Computerized analysis of plantar pressure variation in flip-flops, athletic shoes, and bare feet [Carl and Barrett], 5:374
Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36
Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112
Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379
Reliability of the modified Foot Posture Index [Cornwall et al], 1:7
Transmissions within the tarsal gearbox [Wolf et al], 1:45
Vertical facial dimensions linked to abnormal foot motion [Rothbart], 3:189

Blood
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Body mass
Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379

Bone
Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107
Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171
First metatarsal callus distraction [Benson and Banks], 1:51
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314
Transmissions within the tarsal gearbox [Wolf et al], 1:45
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Tuberculosis of the midtarsal joints: a case report [Jerome], 3:246
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422

Book reviews
Diabetic Neuropathy: Clinical Management, 2nd Ed (Veves and Malik, eds.) [Wu], 3:250

Botulinum toxin type A
Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156
Nonsurgical treatment for hallux abducto valgus with botulinum toxin A [Radovic and Shah], 1:61

Brostrom procedure
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473

Bunionectomy
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

C

Calcaneus
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422

Callus
First metatarsal callus distraction [Benson and Banks], 1:51
What's your diagnosis? [Lemont], 2:137, 6:477

Cancer
Distant metastases to the hallux in nasopharyngeal carcinoma [Lee], 3:239
Prostate cancer metastases to the leg, ankle, and foot [Lee], 3:242

Cantharidin
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445

Carcinoma
Distant metastases to the hallux in nasopharyngeal carcinoma [Lee], 3:239
Prostate cancer metastases to the leg, ankle, and foot [Lee], 3:242

Cardiovascular disease
Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489

Cartilage
Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318

Casting
Ability of foot care professionals to cast feet using the non-weightbearing plaster and the gait-referenced foam casting techniques [Trotter and Pierrynowski], 1:14
Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457

Cephalosporin
Penicillin and cephalosporin drug allergies: a paradigm shift [Smith], 6:479

Charcot's foot
Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1

Chondromatosis
Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318

Claw toe
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27

Clubfoot
Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408
Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451
Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Compartment syndrome
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404

Congenital deformity
Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408
Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451
Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Contracture, ischemic
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404

Corn
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345
What's your diagnosis? [Lemont], 2:137

Costs, health-care
The right to bear legs—an amendment to healthcare: how preventing amputations can save billions for the US health-care system (letter) [Rogers et al], 2:166

Cycling
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261

D

Depression
The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130

Dermatitis
Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42

Dermatophytes
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353

Desquamation
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Diabetes mellitus
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1
Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489
Negative-pressure wound therapy and diabetic foot amputations: a retrospective study of payer claims data (letter) [Udell], 2:164; (author's response) [Frykberg and Williams], 2:165
Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322
Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229
The right to bear legs—an amendment to healthcare: how preventing amputations can save billions for the US health-care system (letter) [Rogers et al], 2:166

Distraction
First metatarsal callus distraction [Benson and Banks], 1:51

E

Education
Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330

Elephantiasis nostras verrucosa
Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66

Equinus deformity
Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Examination, physical
Reliability and validity of current physical examination techniques of the foot and ankle [Wrobel and Armstrong], 3:197

Exercise
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107

Extracorporeal shockwave treatment
Extracorporeal shockwave therapy for the treatment of Achilles tendinopathies: a prospective study [Fridman et al], 6:466

F

Fascia, plantar
Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379

Flatfoot
The flat-footed child—to treat or not to treat: what is the clinician to do? [Evans], 5:386

Foot Posture Index
Reliability of the modified Foot Posture Index [Cornwall et al], 1:7

Foot structure
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112

Foot type
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112
Vertical facial dimensions linked to abnormal foot motion [Rothbart], 3:189

Footwear
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Computerized analysis of plantar pressure variation in flip-flops, athletic shoes, and bare feet [Carl and Barrett], 5:374
Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229
The safety and efficacy of Pain Checker socks in the treatment of mild-to-moderate foot pain: a clinical trial [Jennings et al], 4:278

Forefoot
Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171

Fracture
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422

Fungus
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353

G

Gait
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302
Changes in gait economy between full-contact custom-made foot orthoses and prefabricated inserts in patients with musculoskeletal pain: a randomized controlled trial [Trotter and Pierrynowski], 6:429
Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268
Computerized analysis of plantar pressure variation in flip-flops, athletic shoes, and bare feet [Carl and Barrett], 5:374
Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112
Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Gearbox, tarsal
Transmissions within the tarsal gearbox [Wolf et al], 1:45

Gel, platelet
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Geriatric population
The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130

Graft
Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139

Granuloma
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Guidelines, authors'
Guidelines for authors [Joseph], 1:79, 4:339

H

Hallux
Distant metastases to the hallux in nasopharyngeal carcinoma [Lee], 3:239
Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Nonsurgical treatment for hallux abducto valgus with botulinum toxin A [Radovic and Shah], 1:61

Hallux abducto valgus
Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Nonsurgical treatment for hallux abducto valgus with botulinum toxin A [Radovic and Shah], 1:61

Hallux limitus
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123

Hallux rigidus
Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

Hammer toe
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27

Heel
Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212

Heloma durum
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Heloma neurofibrosum
What's your diagnosis? [Lemont], 2:137

Hyperhidrosis
Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156

Hyperparathyroidism
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

I

Imaging
Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171
Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Surgical correlation of preoperative MRI findings of trauma to tendons and ligaments of the foot and ankle [Kuwada], 5:370

Infection
Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1
Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
Tuberculosis of the midtarsal joints: a case report [Jerome], 3:246
Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139

Injury
Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404
The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212
Surgical correlation of preoperative MRI findings of trauma to tendons and ligaments of the foot and ankle [Kuwada], 5:370
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422
Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

Insole
Changes in gait economy between full-contact custom-made foot orthoses and prefabricated inserts in patients with musculoskeletal pain: a randomized controlled trial [Trotter and Pierrynowski], 6:429
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36
Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229
The short-term effectiveness of full-contact custom-made foot orthoses and prefabricated shoe inserts on lower-extremity musculoskeletal pain: a randomized clinical trial [Trotter and Pierrynowski], 5:357

Intertrigo
Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42

Ischemia
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404

J

Joint
Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
Clinical and radiographic features of pigmented villonodular synovitis of the foot: report of eight cases (letter) [Saxena], 3:251; (author's response) [Carpintero], 3:251
Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1
Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Nonsurgical treatment for hallux abducto valgus with botulinum toxin A [Radovic and Shah], 1:61
The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207
Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379
Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27
Transmissions within the tarsal gearbox [Wolf et al], 1:45
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Tuberculosis of the midtarsal joints: a case report [Jerome], 3:246
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

K

Keratosis
What's your diagnosis? [Lemont], 6:477

Kinematics
Transmissions within the tarsal gearbox [Wolf et al], 1:45

Kirschner wire
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

Knee
The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207

L

Lesion
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
Clinical and radiographic features of pigmented villonodular synovitis of the foot: report of eight cases (letter) [Saxena], 3:251; (author's response) [Carpintero], 3:251
The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
Distant metastases to the hallux in nasopharyngeal carcinoma [Lee], 3:239
Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66
Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
Prostate cancer metastases to the leg, ankle, and foot [Lee], 3:242
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345
Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314
Tuberculosis of the midtarsal joints: a case report [Jerome], 3:246
Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311
What's your diagnosis? [Lemont], 2:137, 4:337, 6:477
Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

Ligament
Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207
Surgical correlation of preoperative MRI findings of trauma to tendons and ligaments of the foot and ankle [Kuwada], 5:370
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473

Lymphedema
A comparison of different assessment techniques for measuring foot and ankle volume in healthy adults [Friends et al], 2:85
Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66

M

Maceration
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353

Matrixectomy
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418

Measurement
Ability of foot care professionals to cast feet using the nonweightbearing plaster and the gait-referenced foam casting techniques [Trotter and Pierrynowski], 1:14
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302
A comparison of different assessment techniques for measuring foot and ankle volume in healthy adults [Friends et al], 2:85
Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268
Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Metatarsus adductus angle in male and female feet: normal values with two measurement techniques [Domínguez and Munuera], 5:364
Reliability and validity of current physical examination techniques of the foot and ankle [Wrobel and Armstrong], 3:197
Reliability of the modified Foot Posture Index [Cornwall et al], 1:7

Meniscus
Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Metatarsal
First metatarsal callus distraction [Benson and Banks], 1:51
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Metatarsus adductus angle in male and female feet: normal values with two measurement techniques [Domínguez and Munuera], 5:364
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Metatarsus adductus angle
Metatarsus adductus angle in male and female feet: normal values with two measurement techniques [Domínguez and Munuera], 5:364

Microscopy
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224

Morphology
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112
The flat-footed child—to treat or not to treat: what is the clinician to do? [Evans], 5:386
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Vertical facial dimensions linked to abnormal foot motion, [Rothbart] 3:189

Muscle
Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408
Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139

N

Nail
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Negative-pressure wound therapy
Negative-pressure wound therapy and diabetic foot amputations: a retrospective study of payer claims data (letter) [Udell], 2:164; (author's response) [Frykberg and Williams], 2:165

Neuroarthropathy
Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1

Neuroischemia
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143

Neuropathy
Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261

Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489
Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322

O

Onychocryptosis
Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Onychomycosis
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224

Orthoses
Ability of foot care professionals to cast feet using the non-weightbearing plaster and the gait-referenced foam casting techniques [Trotter and Pierrynowski], 1:14
Changes in gait economy between full-contact custom-made foot orthoses and prefabricated inserts in patients with musculoskeletal pain: a randomized controlled trial [Trotter and Pierrynowski], 6:429
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
A comparison of rearfoot motion control and comfort between custom and semicustom foot orthotic devices [Davis et al], 5:394
Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36
The flat-footed child—to treat or not to treat: what is the clinician to do? [Evans], 5:386
Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229
The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207
The short-term effectiveness of full-contact custom-made foot orthoses and prefabricated shoe inserts on lower-extremity musculoskeletal pain: a randomized clinical trial [Trotter and Pierrynowski], 5:357

Osteochondromatosis
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70

Osteogenesis
First metatarsal callus distraction [Benson and Banks], 1:51

Osteolysis
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1

Osteomyelitis
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1
Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139

P

Pain
Changes in gait economy between full-contact custom-made foot orthoses and prefabricated inserts in patients with musculoskeletal pain: a randomized controlled trial [Trotter and Pierrynowski], 6:429
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107
Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283
The safety and efficacy of Pain Checker socks in the treatment of mild-to-moderate foot pain: a clinical trial [Jennings et al], 4:278
The short-term effectiveness of full-contact custom-made foot orthoses and prefabricated shoe inserts on lower-extremity musculoskeletal pain: a randomized clinical trial [Trotter and Pierrynowski], 5:357

Pediatric population
Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268
Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408
Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451
The flat-footed child—to treat or not to treat: what is the clinician to do? [Evans], 5:386
Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Penicillin
Penicillin and cephalosporin drug allergies: a paradigm shift [Smith], 6:479

Peripheral arterial disease
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143

Pes planus
The flat-footed child—to treat or not to treat: what is the clinician to do? [Evans], 5:386

Pharmacology
Penicillin and cephalosporin drug allergies: a paradigm shift, 6:479

Phenol
Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418

Plasminogen activator
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Platelet
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Podophyllotoxin
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445

Polymerase chain reaction
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224

Porokeratosis plantaris discreta (Steinberg's lesion)
What's your diagnosis? [Lemont], 4:337

President's Message
Acceptance address [Taubman], 3:254
APMA PAC [Robertozzi], 1:84
Board certification and parity [Taubman], 6:495
Finding the future, being a mentor, and parity [Taubman], 5:427
Parity and residency training and development [Taubman], 4:342
Public health and podiatry [Robertozzi], 2:169
Thank you [Robertozzi], 3:253

Pressure, plantar
Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143
Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268
Computerized analysis of plantar pressure variation in flip-flops, athletic shoes, and bare feet [Carl and Barrett], 5:374
Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112
Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229

Profession, podiatric medical
Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330

Pronation
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379

Prostate
Prostate cancer metastases to the leg, ankle, and foot [Lee], 3:242

Prosthesis
Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Protein, C-reactive
Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot [Judge], 1:1

Psychology
The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130

Public health
Public health and podiatry (letter) [Levy], 5:426; (author's response) [Robertozzi], 5:426

Q

Quality of life
Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283

R

Rearfoot
A comparison of rearfoot motion control and comfort between custom and semicustom foot orthotic devices [Davis et al], 5:394

Running
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36

S

Salicylic acid
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Sesamoid
Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149

Sever's disease
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212

Shin splints
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107

Skin
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66
Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353
Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345
What's your diagnosis? [Lemont], 2:137, 4:337, 6:477

Sports medicine
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107
The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207
Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212

Sprain
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473

Steinberg's lesion
What's your diagnosis? [Lemont], 4:337

Surgery
Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118
Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451
First metatarsal callus distraction [Benson and Banks], 1:51
Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418
Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296
Surgical correlation of preoperative MRI findings of trauma to tendons and ligaments of the foot and ankle [Kuwada], 5:370
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422
Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

Syndesmosis
Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469

Synovitis
Clinical and radiographic features of pigmented villonodular synovitis of the foot: report of eight cases (letter) [Saxena], 3:251; (author's response) [Carpintero], 3:251

Synovium
Clinical and radiographic features of pigmented villonodular synovitis of the foot: report of eight cases (letter) [Saxena], 3:251; (author's response) [Carpintero], 3:251
Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70

T

Talus
Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473

Tarsus
Transmissions within the tarsal gearbox [Wolf et al], 1:45
Tuberculosis of the midtarsal joints: a case report [Jerome], 3:246
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422

Tendon
Extracorporeal shockwave therapy for the treatment of Achilles tendinopathies: a prospective study [Fridman et al], 6:466
Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414
Surgical correlation of preoperative MRI findings of trauma to tendons and ligaments of the foot and ankle [Kuwada], 5:370
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27
What's your diagnosis? [Lemont], 6:477
Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

Tibia
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107

Tinea pedis
The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353

Toe
Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418
Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296
Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Tuberculosis
Tuberculosis of the midtarsal joints: a case report [Jerome], 3:246

Tumor
Distant metastases to the hallux in nasopharyngeal carcinoma [Lee], 3:239
Prostate cancer metastases to the leg, ankle, and foot [Lee], 3:242
Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

U

Ulcer
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489
Negative-pressure wound therapy and diabetic foot amputations: a retrospective study of payer claims data (letter) [Udell], 2:164; (author's response) [Frykberg and Williams], 2:165
Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
The right to bear legs—an amendment to healthcare: how preventing amputations can save billions for the US health-care system (letter) [Rogers et al], 2:166
Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures [Hollawell], 5:422
Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

V

Vamp disease
What's your diagnosis? [Lemont], 6:477

Vascular system
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489
Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322

Verruca
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
What's your diagnosis? [Lemont], 2:137

Virus
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
What's your diagnosis? [Lemont], 2:137

Vitamin D
Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Volume, limb
A comparison of different assessment techniques for measuring foot and ankle volume in healthy adults [Friends et al], 2:85

W

Wart
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
What's your diagnosis? [Lemont], 2:137

Workforce
Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330

AUTHOR INDEX

A

Agus, Haluk: Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451

Aksoy, Murat: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290

Alarcón García, Juan Maria: Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379

Altay, Taskin: Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408

Angirasa, Arush K.: Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473

Armstrong, David G.:

Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153

CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261

Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322

Reliability and validity of current physical examination techniques of the foot and ankle [Wrobel and Armstrong], 3:197

The right to bear legs—an amendment to healthcare: how preventing amputations can save billions for the US health-care system (letter) [Rogers et al], 2:166

Aron, David: Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489

Atway, Said: Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171

Augustine, Elizabeth: A comparison of different assessment techniques for measuring foot and ankle volume in healthy adults [Friends et al], 2:85

Avil, Steven J.:

Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436

Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107

Aynaci, Osman: Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318

B

Baba, Fusun: Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Baktiroglu, Selcuk: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290

Banks, Alan S.: First metatarsal callus distraction [Benson and Banks], 1:51

Barnes, Mike R.: Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107

Barrett, Michael J.: Talar anchor placement for modified Brostrom lateral ankle stabilization procedure [Angirasa and Barrett], 6:473

Barrett, Stephen L.: Computerized analysis of plantar pressure variation in flip-flops, athletic shoes, and bare feet [Carl and Barrett], 5:374

Becerro de Bengoa Vallejo, Ricardo:

Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445

Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418

Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27

Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

Begg, Lindy: Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143

Bell, Bryan L.: Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70

Belpedio, Drew: Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311

Benedetti, Maria Grazia: Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Benson, J. Christopher: First metatarsal callus distraction [Benson and Banks], 1:51

Berberoglu, Kezban: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290

Berti, Lisa: Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Bevilacqua, Nicholas J.:

Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153

Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322

Blau, Jill C.: Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156

Boesiger, Peter: Transmissions within the tarsal gearbox [Wolf et al], 1:45

Burns, Joshua:

Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143

Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112

Butler, Robert J.: Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102

C

Cain, Jarrett: Extracorporeal shockwave therapy for the treatment of Achilles tendinopathies: a prospective study [Fridman et al], 6:466

Calangu, Semra: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290

Camp Faulí, Angel: Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229
Campbell, Jackie A.:
Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436
Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107
Carl, Tanya J.: Computerized analysis of plantar pressure variation in flip-flops, athletic shoes, and bare feet [Carl and Barrett], 5:374
Carpintero, Pedro: Clinical and radiographic features of pigmented villonodular synovitis of the foot: report of eight cases (author's response), 3:251
Catani, Fabio: Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19
Catanzariti, Alan R.:
Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Cetinkaya, Zafer: Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42
Child, Brandon: Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Chipchase, Lucy S.: Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Christina, James R.: Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330
Christodoulou, Anastasios: Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160
Çobanoglu, Ümit: Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318
Cook, Jill L.: Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283
Córdoba-Fernandez, Antonio: Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296
Cornwall, Mark W.: Reliability of the modified Foot Posture Index [Cornwall et al], 1:7
Cosin Matamoros, Eva: Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379
Cosin Matamoros, Julia: Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379
Crews, Ryan T.: CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Cromack, Douglas T.: Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Crosbie, Jack: Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112

D

Danoff, Jerome: A comparison of different assessment techniques for measuring foot and ankle volume in healthy adults [Friends et al], 2:85
Davis, Irene S.:
Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102
A comparison of rearfoot motion control and comfort between custom and semicustom foot orthotic devices [Davis et al], 5:394
DeHeer, Patrick: Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66
DeLeo, Alison T.: A comparison of rearfoot motion control and comfort between custom and semicustom foot orthotic devices [Davis et al], 5:394
DellaCorte, Michael P.: Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153
Demiralp, Bahtiyar: Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Demirel, Reha: Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42
DeMore, Matthew, III: Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171
Desideri, Carla Maria: The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
Díaz Martínez, Teresa: Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379
Dionne, Mark G.: Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330
Dogan, Ali: Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414
Domínguez, Gabriel:
Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Metatarsus adductus angle in male and female feet: normal values with two measurement techniques [Domínguez and Munuera], 5:364
Duckworth, Amy L.: Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66
Dunn, S. Patrick: Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156

E

Erler, Kaan: Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Ertugrul, M. Bulent: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290
Evans, Angela Margaret: The flat-footed child—to treat or not to treat: what is the clinician to do? 5:386

F

Fernández, Maria José: Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229

Forte, Gaetano J.: Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330

Francis, Kenneth: Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153

Fridman, Robert: Extracorporeal shockwave therapy for the treatment of Achilles tendinopathies: a prospective study [Fridman et al], 6:466

Friends, John: A comparison of different assessment techniques for measuring foot and ankle volume in healthy adults [Friends et al], 2:85

Frykberg, Robert G.: Negative-pressure wound therapy and diabetic foot amputations: a retrospective study of payer claims data (author's response), 2:164

G

Gauthier, Caroline: Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156

Giannini, Sandro: Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Giri, Subhasis K.: Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118

Givissis, Panagiotis: Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160

Gómez-Martín, Beatriz: Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445

Goodwin, Susan: The safety and efficacy of Pain Checker socks in the treatment of mild-to-moderate foot pain: a clinical trial [Jennings et al], 4:278

Gracely, Edward: Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268

Grady, John F.: The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353

Gupta, Aditya K.: Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224

H

Heda, Ghanshyam D.: Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Heiderscheit, Bryan: Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36

Hernández-Cortés, Pedro: Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404

Hernández-Hernández, Miguel Angel: Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404

Hillstrom, Howard: Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102

Hollawell, Shane: Wound closure technique for lateral extensile approach to intra-articular calcaneal fractures, 5:422

Hunt, Adrienne: Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112

Husain, Jugnoo: Elephantiasis nostras verrucosa or "mossy foot lesions" in lymphedema praecox: report of a case [Duckworth et al], 1:66

I

Irving, Damien B.: Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283

J

Jafri, Mansoor: Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118

Jenkins, Walter L.: The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207

Jennings, Maureen B.: The safety and efficacy of Pain Checker socks in the treatment of mild-to-moderate foot pain: a clinical trial [Jennings et al], 4:278

Jerome, J. Terrence Jose: Tuberculosis of the midtarsal joints: a case report, 3:246

John, Shine: Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75

Jones, Sara: Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212

Joseph, Warren S.: Guidelines for authors, 1:79, 4:339

Juarez-Jiménez, José María: Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Judge, Molly S.: Infection and neuroarthropathy: the utility of C-reactive protein as a screening tool in the Charcot foot, 1:1

Jules, Kevin T.:

Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418

Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27

Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

K

Kalender, Ali Murat: Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Kalenderer, Onder: Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451

Kaplan, Yelena: Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489

Karaca, Semsettin: Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42

Karapinar, Levent: Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408

Kaya, Ahmet: Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408
Keane, Ralph: Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118
Kerimoglu, Servet: Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318
Klein, Erin E.: CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Komurcu, Mahmut: Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469
Kuba, Jennifer: Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489
Kulac, Mustafa: Etiology of foot intertrigo in the district of Afyonkarahisar, Turkey: a bacteriologic and mycologic study [Karaca et al], 1:42
Kuwada, Gerald T.: Surgical correlation of preoperative MRI findings of trauma to tendons and ligaments of the foot and ankle, 5:370

L

Lacey, Autum M.: The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353
Lafuente, Guillermo: Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123
Lamm, Bradley M.: Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Lange, Belinda S.: Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Lavery, Lawrence A.: The right to bear legs—an amendment to healthcare: how preventing amputations can save billions for the US health-care system (letter) [Rogers et al], 2:166
Leardini, Alberto: Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19
Lebec, Michael: Reliability of the modified Foot Posture Index [Cornwall et al], 1:7
Lee, Daniel K.:
Distant metastases to the hallux in nasopharyngeal carcinoma, 3:239
Prostate cancer metastases to the leg, ankle, and foot, 3:242
Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70
Legge, Bradford S.: The incidence of tinea pedis in diabetic versus nondiabetic patients with interdigital macerations: a prospective study [Legge et al], 5:353
Lemont, Harvey: What's your diagnosis? 2:137, 4:337, 6:477
Lesczczynski, Christine: The safety and efficacy of Pain Checker socks in the treatment of mild-to-moderate foot pain: a clinical trial [Jennings et al], 4:278
Levy, Leonard A.: Public health and podiatry (letter), 5:426
Li, Albert: Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95
Llobell Andrés, Cristina: Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229
Long, Jodi: Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489
Longo, Rosella: The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
Losa Iglesias, Marta Elena:
Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445
Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326
Louk, Louis, Jr.: Synovial osteochondromatosis involvement in post-traumatic ankle injury [Lee et al], 1:70
Luechinger, Roger: Transmissions within the tarsal gearbox [Wolf et al], 1:45

M

Maffulli, Nicola: Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95
Malik, Rayaz A.: Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322
Mannucci, Edoardo: The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
Marchionni, Niccolò: The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
Martínez-Nova, Alfonso: Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302
Maskill, Michael: Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Masotti, Giulio: The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130
McClellan, Scott: Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311
McPoil, Thomas G.: Reliability of the modified Foot Posture Index [Cornwall et al], 1:7
Mendicino, Robert W.:
Long leg calcaneal axial and hindfoot alignment radiographic views for frontal plane assessment [Mendicino et al], 1:75
Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149
Menz, Hylton B.: Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283
Monami, Matteo: The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms [Monami et al], 2:130

Montiel Parreño, Enrique: Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229

Munuera, Pedro V.:

Length of the sesamoids and their distance from the metatarsophalangeal joint space in feet with incipient hallux limitus [Munuera et al], 2:123

Metatarsus adductus angle in male and female feet: normal values with two measurement techniques [Domínguez and Munuera], 5:364

N

Nguyen, Aidan: Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

O

Oladeji, Oluwabunmi: Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268

O'Leary, Katherine: Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36

Orgilés Barceló, César: Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229

Orhan, Ismahan: Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Orlin, Margo: Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268

Orphanos, Joann: Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139

Osher, Lawrence S.: Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171

Ozturan, Kutay: Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469

Ozturk, Adil: Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Ozturk, Cagatay: Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469

Ozturk, Hasan: Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408

P

Pajares-López, Miguel: Ischemic contracture of deep posterior compartment of the leg following isolated ankle fracture [Hernández-Cortés et al], 5:404

Pascual Huerta, Javier:

Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302

Relationship of body mass index, ankle dorsiflexion, and foot pronation on plantar fascia thickness in healthy, asymptomatic subjects [Pascual Huerta et al], 5:379

Pepe, Linda: Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311

Pierrynowski, Michael Raymond:

Ability of foot care professionals to cast feet using the non-weightbearing plaster and the gait-referenced foam casting techniques [Trotter and Pierrynowski], 1:14

Changes in gait economy between full-contact custom-made foot orthoses and prefabricated inserts in patients with musculoskeletal pain: a randomized controlled trial [Trotter and Pierrynowski], 6:429

The short-term effectiveness of full-contact custom-made foot orthoses and prefabricated shoe inserts on lower-extremity musculoskeletal pain: a randomized clinical trial [Trotter and Pierrynowski], 5:357

Porta Rosas, Norberto: Physical evaluation of insole materials used to treat the diabetic foot [Camp Faulí et al], 3:229

Pournaras, John: Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160

Prados Frutos, Juan Carlos:

Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27

Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326

R

Radovic, Philip A.: Nonsurgical treatment for hallux abducto valgus with botulinum toxin A [Radovic and Shah], 1:61

Raedeke, Susanne G.: The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207

Rayo-Rosado, Rafael: Platelet gel for the surgical treatment of onychocryptosis [Córdoba-Fernandez et al], 4:296

Reisoglu, Ali: Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451

Richards, Christine J.: Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102

Robbins, Jeffrey M.: Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489

Robertozzi, Christian A.:

APMA PAC, 1:84

Public health and podiatry, 2:169; (author's response), 5:426

Thank you, 3:253

Roberts, Lee K.: Role of tissue-type plasminogen activator in salicylic acid–induced sloughing of human corn tissue [Heda and Roberts], 5:345

Rogers, Lee C.:

Charcot's arthropathy in a patient with HIV-associated neuropathy [Rogers et al], 2:153

Neuropathy in diabetes: not a knee-jerk diagnosis [Rogers et al], 4:322

The right to bear legs—an amendment to healthcare: how preventing amputations can save billions for the US healthcare system (letter) [Rogers et al], 2:166

Romagnoli, Matteo: Functional outcome of meniscal-bearing total ankle replacement: a gait analysis study [Benedetti et al], 1:19

Rothbart, Brian A.: Vertical facial dimensions linked to abnormal foot motion, 3:189

Roukis, Thomas S.: Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139

S

Sáez Crespo, Antonio: Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445

Salman, Serpil: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290

Saltrick, Karl: Traumatic dislocation of the first metatarsophalangeal joint with tibial sesamoid fracture: a case report of a type III B dislocation [Maskill et al], 2:149

Sánchez Gómez, Rubén:

Application of cantharidin and podophyllotoxin for the treatment of plantar warts [Becerro de Bengoa Vallejo et al], 6:445

Gauze application of phenol for matrixectomy [Becerro de Bengoa Vallejo et al], 5:418

Sánchez-Rodríguez, Raquel: Cadence, age, and weight as determinants of forefoot plantar pressures using the Biofoot in-shoe system [Martínez-Nova et al], 4:302

Saraçoglu, Metehan: Synovial chondromatosis of the subtalar joint: a case report and review of the literature [Kerimoglu et al], 4:318

Saxena, Amol:

Clinical and radiographic features of pigmented villonodular synovitis of the foot: report of eight cases (letter), 3:251

Wound complications from surgeries pertaining to the Achilles tendon: an analysis of 219 surgeries [Saxena et al], 2:95

Scharfbillig, Rolf W.: Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212

Scutter, Sheila D.: Sever's disease: what does the literature really tell us? [Scharfbillig et al], 3:212

Sebik, Ahmet: Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Seramet, Ebubekir: Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

Shah, Ekta: Nonsurgical treatment for hallux abducto valgus with botulinum toxin A [Radovic and Shah], 1:61

Shaikh, Faisal M.: Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails [Shaikh et al], 2:118

Sharrer, Edward: Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311

Singh, Jagpal: Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224

Sirmatel, Ocal: Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314

Smith, Robert G.: Penicillin and cephalosporin drug allergies: a paradigm shift, 6:479

Song, Jinsup: Arch height index measurement system: establishment of reliability and normative values [Butler et al], 2:102

Stackhouse, Carrie: Comparison of the two-step and midgait methods of plantar pressure measurement in children [Oladeji et al], 4:268

Stacoff, Alex: Transmissions within the tarsal gearbox [Wolf et al], 1:45

Strauss, Gerald: Mortality rates and diabetic foot ulcers: is it time to communicate mortality risk to patients with diabetic foot ulceration? [Robbins et al], 6:489

Strobel, Stephen: Unusual infection in the foot of a barefoot gardener [Strobel et al], 4:311

Stuessi, Edgar: Transmissions within the tarsal gearbox [Wolf et al], 1:45

Surenkok, Fatih: Congenital clubfoot associated with an accessory soleus muscle [Karapinar et al], 5:408

Symeonidis, Panagiotis: Interposition arthroplasty of the first metatarsophalangeal joint with a fascia lata allograft [Givissis et al], 2:160

T

Tasatan, Ersin: Acute traumatic open posterolateral dislocation of the ankle without tearing of the tibiofibular syndesmosis ligaments: a case report [Demiralp et al], 6:469

Taubman, Ross E.:

Acceptance address, 3:254

Board certification and parity, 6:495

Finding the future, being a mentor, and parity, 5:427

Parity and residency training and development, 4:342

Thomson, Matthew K.: Extended pedal imaging via modifications of the traditional forefoot axial radiographic study: teaching new tricks to an old dog? an initial report with case presentations [Osher et al], 3:171

Trotter, Leslie C.:

Ability of foot care professionals to cast feet using the nonweightbearing plaster and the gait-referenced foam casting techniques [Trotter and Pierrynowski], 1:14

Changes in gait economy between full-contact custom-made foot orthoses and prefabricated inserts in patients with musculoskeletal pain: a randomized controlled trial [Trotter and Pierrynowski], 6:429

The short-term effectiveness of full-contact custom-made foot orthoses and prefabricated shoe inserts on lower-extremity musculoskeletal pain: a randomized clinical trial [Trotter and Pierrynowski], 5:357

Turgut, Ali: Evaluation of clinical and radiographic outcomes of complete subtalar release in clubfoot treatment [Kalenderer et al], 6:451

Tweed, Jo L.:

Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners [Tweed et al], 6:436

Etiologic factors in the development of medial tibial stress syndrome: a review of the literature [Tweed et al], 2:107

U

Udell, Elliot: Negative-pressure wound therapy and diabetic foot amputations: a retrospective study of payer claims data (letter), 2:164

Unal, Seher: Pathogens isolated from deep soft tissue and bone in patients with diabetic foot infections [Ertugrul et al], 4:290

Uslu, Mustafa: Mini-open technique for the Achilles tenotomy in correction of idiopathic clubfoot: a report of 25 cases [Dogan et al], 5:414

V

Vicaretti, Mauro: Comparison of orthotic materials on foot pain, comfort, and plantar pressure in the neuroischemic diabetic foot: a case report [Burns et al], 2:143

Vicenzino, Bill: Reliability of the modified Foot Posture Index [Cornwall et al], 1:7

Viejo Tirado, Fermín:
Transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 1:27
Use of a Kirschner wire for distraction and capsular flaps in the Keller interpositional arthroplasty [Becerro de Bengoa Vallejo et al], 4:326
Vlahovic, Tracey C.: Injectable botulinum toxin as a treatment for plantar hyperhidrosis: a case study [Vlahovic et al], 2:156
Vorpahl, Kristin Anderson: Effect of cushioned insoles on impact forces during running [O'Leary et al], 1:36

W

Walters, Julie: Effect of a low-Dye application of Scotchcast soft cast on peak and mean plantar pressures in subjects with a navicular drop greater than 10 mm [Lange et al], 6:457
Weil, Lowell, Jr.: Extracorporeal shockwave therapy for the treatment of Achilles tendinopathies: a prospective study [Fridman et al], 6:466
Weil, Lowell, Sr.: Extracorporeal shockwave therapy for the treatment of Achilles tendinopathies: a prospective study [Fridman et al], 6:466
Williams, D.S. Blaise, III: The relationship between the use of foot orthoses and knee ligament injury in female collegiate basketball players [Jenkins et al], 3:207
Williams, David V.: Negative-pressure wound therapy and diabetic foot amputations: a retrospective study of payer claims data (author's response), 2:164
Wilson, Jodi: Reliability of the modified Foot Posture Index [Cornwall et al], 1:7
Wing, Paul: Projections of the supply of and demand for podiatric physicians in the United States, 2005 to 2030 [Wing et al], 4:330
Wolf, Peter: Transmissions within the tarsal gearbox [Wolf et al], 1:45
Wong, Lilian: Effect of foot morphology on center-of-pressure excursion during barefoot walking [Wong et al], 2:112
Wrobel, James S.:
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Reliability and validity of current physical examination techniques of the foot and ankle [Wrobel and Armstrong], 3:197
Wu, Stephanie C.:
CLEAR Cleat: a proof-of-concept trial of an aerobic activity facilitator to reduce plantar forefoot pressures and their potential in those with foot ulcers [Klein et al], 4:261
Review of *Diabetic Neuropathy: Clinical Management*, 2nd Ed (Veves and Malik, eds), 3:250

Y

Yazgan, Pelin: Third metatarsal brown tumor with secondary hyperparathyroidism: an atypical localization [Yazgan et al], 4:314
Young, Mark A.: Impact of chronic plantar heel pain on health-related quality of life [Irving et al], 4:283

Z

Zaman, Muhammad: Diagnosis of *Trichophyton rubrum* from onychomycotic nail samples using polymerase chain reaction and calcofluor white microscopy [Gupta et al], 3:224
Zgonis, Thomas: Use of a muscle flap and a split-thickness skin graft for a calcaneal osteomyelitis after an open reduction and internal fixation [Zgonis et al], 2:139
Zifchock, Rebecca Avrin: A comparison of rearfoot motion control and comfort between custom and semicustom foot orthotic devices [Davis et al], 5:394

Receive Customized Researcher Alerts by E-Mail

Track new published information that is relevant to your research through JAPMA Online's CiteTrack feature. You can be automatically notified by e-mail when information you track is published in JAPMA or any other participating HighWire-hosted journal. You can choose to receive alerts when a new issue and table of contents are released, when an important article is cited, or when key terms or authors you select appear in new content. This service is free to both subscribers and nonsubscribers. Sign up from any search results page or the Subscriptions page on JAPMA Online at http://www.japmaonline.org/subscriptions.